

HIV/AIDS Awareness, Attitudes and Risk Behavior Among University Students in Wuhan, China

Madelene Albrektsson¹, Louise Alm¹, Xiaodong Tan² and Rune Andersson^{*,1,3}

¹Department of Infectious Diseases, Institute of Biomedicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

²School of Public Health, Wuhan University, Wuhan, China

³Research and Development Centre, Skaraborg Hospital, Skövde, Sweden

Abstract: *Background:* In China, the estimated number of HIV positives in 2007 was 700,000 and the epidemic continues to expand. Because of their attitudes towards sexual behavior, young people are considered to be a group at high risk.

Methods: Eight hundred sixty-eight undergraduate students at Wuhan University were selected through stratified cluster sampling, to answer a questionnaire. They were divided into three main groups: Chinese medical students, foreign medical students and Chinese students from other faculties. Fourteen interviews were conducted in addition.

Results: Ninety-nine percent of the students had heard of HIV/AIDS and 76% of the students could distinguish HIV from AIDS. The main route of transmission was believed by the Chinese students to be blood transfusion and sexual intercourse by the foreign medical students. The female students knew more about the routes of transmission than the male students. Medical students had a higher level of knowledge than non-medical students, and among the medical students, the foreign students were more knowledgeable than the Chinese students. Only 8 % of the students were sexually active.

Conclusion: The students had an accepting attitude towards people living with HIV and no extensive risk behavior. Overall, the knowledge level was found to be moderate.

Keywords: HIV, AIDS, information, epidemic, ethnic.

INTRODUCTION

HIV and AIDS in China

The first case of HIV in China was reported in 1985. The first epidemic spread began in 1989 among injecting drug users in rural areas. Since 1998 all 31 provinces, autonomous regions and municipalities have reported cases of HIV [1]. At the end of October 2007 the reported number of HIV positives was 223,501; including 62,838 cases of AIDS, while by the end of 2007 the estimated number of HIV positives was 700,000 (range 550,000-850,000). This means that the HIV prevalence in China is 0.05 per cent. In comparison with other countries, the overall prevalence rate in China is still low but the epidemic continues to expand. The pace of the expansion has begun to slow down. In 2007, the estimated number of new infections was 50,000 [2]. According to calculations made by UNAIDS in 2000, the number of people infected in China might be as high as 10 million by 2010 [3] The Chinese government has taken actions to control the spread; the goal is to limit the number of HIV cases to 1.5 million in 2010 [4].

During the last few years the main route of transmission has changed from injecting drug users to heterosexual

transmission [2,5]. A contributing factor to the increased heterosexual transmission is a change of attitude towards sexuality in China. Since the declaration of the One Child Family policy, premarital sex has become widely accepted among young people. Sexual behavior is now considered an individual responsibility [5].

Stigmatization of HIV and AIDS is a serious problem throughout China. Many people living with HIV/AIDS are afraid to tell anyone about their status. If they choose to tell people in their surroundings they risk losing their social support and jobs and being discriminated against in other ways [6].

HIV and AIDS in Hubei

Hubei province is in central China, with a population of 65 million. The majority of the people live in rural areas. Together with Yunnan, Guangxi, Anhui and Guangdong, Hubei accounts for 83 per cent of the total reported number of HIV cases in the country [2]. Hubei province is estimated to have around 45,000 HIV positives. This high number is explained by commercial blood selling several years ago. The epidemic started among the poor population in rural areas. Transmission owing to blood selling is now under control but the virus keeps spreading through sexual relations and mother-to-child transmission [6]. The epidemic in Hubei, as elsewhere in China, has spread from high risk groups to the general population [2].

*Address correspondence to this author at the Research and Development Centre, Skaraborg Hospital, SE-541 85 Skövde, Sweden; Tel: +46 500-431137; Fax: +46 500 431121; E-mail: rune.andersson@vregion.se

Wuhan University

Wuhan University, in the capital of Hubei, is a key university directly under the administration of the Education Ministry of the People's Republic of China. The university was founded in 1893 and has 29 colleges and various branches of learning. The number of students enrolled is 53,000, 18,000 of whom are postgraduates [7]. Students come to attend the university from all over China and from many foreign countries.

Objective

The HIV virus is spreading rapidly throughout China in the general population. Young people are considered to be a group at high risk. Therefore it is essential to assess their knowledge, attitudes and risk behavior when planning preventive strategies.

PARTICIPANTS AND METHODS

Sampling

Undergraduate students at Wuhan University were selected, using stratified cluster sampling, to answer a questionnaire during spring 2008. The participants studied at different faculties, including Medicine, Information Science, Engineering, Social Science, Humaniores Science and at the Faculty of Science. The three main groups are presented in Table 1.

Table 1. The 868 Participating Students by University Year

Year	Chinese Medical Students		Chinese Non-Medical Students		Foreign Medical Students	
	Number	%	Number	%	Number	%
First	120	45.3	45	13.0	89	34.9
Second	70	26.4	127	36.7	82	32.2
Third	70	26.4	164	47.4	81	31.8
Fourth	1	0.4	10	2.9	3	1.2
Fifth	4	1.5				

All participating students were informed of the general objectives of the study and that they could choose to withdraw at any time. In order to ensure the confidentiality of the information provided, they were advised not to write their names on the questionnaires.

Questionnaire

To assess the knowledge, attitudes and risk behaviors regarding HIV and AIDS, a self-administrated questionnaire was used. The questionnaire consisted of 47 questions divided into four categories: General questions, Questions about HIV/AIDS, Private life and Questions about work situation. The questions were in English but with some of the words translated into Chinese to clarify. To avoid linguistic misunderstandings, the Chinese students were able to direct their questions to both English and Chinese speaking students during the completion of the form. The time

required to fill out the questionnaire was approximately 20-30 minutes.

The questionnaire used in the study has been previously tested in other international studies [8,9]. Modifications were made to adjust the questionnaire to Chinese culture and customs, such as religious and ethnical groups.

Interviews

For deeper understanding, 14 interviews were conducted. Eight of them were with Chinese students and six were with foreign students. The students attended different departments at the university, 8 were men and 6 were women. The age range was 19 to 45.

All students were informed that their participation was voluntary and that they did not have to answer questions they found too private. The Chinese students could choose to do the interview in Chinese or in English. All interviews with the foreign students were conducted in English.

The interview questions were based on interviews performed in similar studies [8,9]. The interviewed student was asked to talk freely about subjects such as general knowledge about HIV/AIDS, attitudes towards infected persons, sexuality and the possibility of a future epidemic in China. Each interview took approximately 30 minutes.

Data Analysis

To enter and analyze the data, Epidata version 3.1 and the Statistical Package for Social Sciences (SPSS) version 13.0 were used. Data obtained were evaluated by frequency and percentage ratios and Chi-square (X^2) test. A p-value of <0.05 was considered significant.

RESULTS

Sample Characteristics

A total of 868 questionnaires were collected. The questionnaires were gathered from three different groups: Chinese medical students (n=265), Chinese non-medical students (n=346) and foreign medical students (n=257); 50.3% were women, 49.2% were men and 0.5% did not give their sex. Ages ranged from 17 to 34 years and the mean age was 20.4.

In relation to the question of ethnicity, 87.4% of the Chinese students answered Han Chinese. The other Chinese students were divided between Tuija (4.1%), Miao (1.0%), Hui (1.5%) and other ethnicity (6.1%). One student did not answer this question. Of the foreign students, 69.3% came from India, 5.2% from Nepal and 3.1% from Pakistan. Most students stated that they had no religion (61.3%). 11.5% were Muslim and 10.0% were Christians.

Among all respondents, 814 (93.8%) were single, 31 (3.6%) lived with a partner, 6 (0.7%) were married, 3 (0.3%) were widowed, 1 (0.1%) was divorced and 4 (0.5%) had other marital status. Nine (1%) students did not answer. Only 7 (0.8%) students answered that they had children.

More than half of the students (59.8%) were majoring in Medicine, 10.9% in Social Science, 8.4% in Humaniores Science, 7.4% in Engineering, 7.0% in Information Science

and 6.5% studied at the Faculty of Science. The non-medical students had generally been studying for longer than the medical students ($p<0.001$), (Table 1).

Sources of HIV/AIDS Information

Of the 868 students, all but 9 had heard of HIV/AIDS, 52.6% stated that they had known about it for five to ten years. The most common source of first information (50.9%) was media, 40.2% had first heard of it in school. There were no significant differences between the three main student groups or gender. Eighty-eight percent answered that they had been given information about HIV/AIDS before entering university and 71.8% had received information at university. Of the foreign medical students, 97.2% had had information before university compared to 85.0% of the Chinese medical students and 89.5% of the Chinese non-medical students ($p<0.003$). Only 74.1% of the foreign students had received information at university, which was significantly fewer than among both the Chinese medical students (88.5%) and the Chinese non-medical students (84.9%) ($p<0.002$).

Fifty-two percent did not feel that the information had been sufficient. In relation to the medical students, non-medical students answered to a larger extent that the received information had not been sufficient (51.7% versus 39.9%, $p<0.02$). The most requested additional information was about current circumstances and spreading tendencies of AIDS all over the world (Table 2). Of the Chinese medical students, 53.5% reported that the information had been sufficient.

General Knowledge of HIV/AIDS

Seven hundred forty-one (95.4%) students answered that there is a difference between HIV and AIDS. When asked to specify, 659 (75.9%) students could distinguish HIV as a virus and AIDS as an infectious disease; 84.5% of the Chinese medical students and 80.2% of the foreign medical students knew of the difference and could specify it correctly. Significantly fewer Chinese non-medical students (66.2%) could make the same distinctions ($p<0.001$).

Religion had no significant influence on the awareness of HIV.

The main route of transmission was considered to be sexual intercourse (40.4%) while 35.5% believed it to be by blood transfusion. However, there was a significant difference of opinion between foreign students and Chinese students ($p<0.001$). More than half of the Chinese students, both medical (53.5%) and non-medical (50.8%), stated blood transfusion as the main route of transmission, while the foreign medical students (81.9%) stated sexual intercourse.

Knowledge regarding transmission routes is presented in Fig. (1). To some of the questions, there were significant differences between the three main groups. Regarding whether kissing is a route of transmission, more Chinese than foreign students believed it to be so (Chinese medical students 29.5%, Chinese non-medical students 32.7% versus foreign medical students 27%, $p<0.01$). The Chinese students believed that sharing needles is a route of transmission to a less extent than foreign students (Chinese medical students 91.6%, Chinese non-medical students 92.4% versus foreign medical students 98.0%, $p<0.04$). Fifty-eight percent of the Chinese non-medical students answered that breastfeeding can spread the disease, which was significantly lower than the Chinese medical students (64.5%) and the foreign medical students (70.4%) ($p<0.04$). Thirty-nine percent of both the Chinese medical students and the Chinese non-medical students believed that the disease can be spread by mosquitoes while only 15.5% of the foreign medical students believed so ($p<0.001$). Regarding legal blood/plasma donation, 38.3% of the Chinese medical students stated this as a route of transmission, which was significantly higher than both the Chinese non-medical students (28.0%) and the foreign medical students (28.9%) ($p<0.03$). Of the female students, 96.3% recognized sharing needles as a route of transmission, compared with 91.1% of the male students ($p<0.01$). Breastfeeding was believed to spread the virus by 69.2% of the women and 57.7% of the men; 66.4% of the women and 56.6% of the men recognised kissing as being harmless and 66.3% of the men and 55.7%

Table 2. Requested Additional Information About HIV/AIDS

Requested information	Chinese Medical Students	Chinese Non-Medical Students	Foreign Medical Students	Total Number	Total %
Current circumstances and spreading tendencies of AIDS all over the world	28	44	32	104	12.0
Prevention-control measures	25	53	22	100	11.5
Research progress	23	29	23	75	8.6
The spreading circumstances of HIV/AIDS in close surroundings	11	20	11	42	4.8
Routes of transmission	4	21	5	30	3.5
Other information	6	2	4	12	1.4
Harmful results	4	4	1	9	1.0
Total	101	173	98	372	42.9
No answer	164	173	159	496	57.1
Total	265	346	257	868	100

Fig. (3). Believed risk for a future HIV/AIDS epidemic in China.

LIMITATIONS

The fact that Wuhan University has a very good academic reputation could be seen as a limitation when the study is compared to studies performed at other universities in China.

The questionnaire was written in English with difficult words translated into Chinese. Still, some of the Chinese students found it difficult to comprehend. To the two questions about whom to talk to about general information of HIV/AIDS and whom to tell if infected, the similar alternatives were unfortunately in a different order in the questionnaire.

DISCUSSION

Sources of HIV/AIDS Information

In the study, it was found that 99% of the students had heard of HIV/AIDS and that the media were the most common sources of information. This coincides with previous studies performed at Wuhan University in 2006 and Xinjiang Medical University in 2004 [8,10]. Only 40.2% reported that their first information was received at school. The Chinese students stated in the interviews that they had had little or no sexual education or information about STD's at school. They all wished that they had been better informed at school and agreed that there is a great need for more sexual education among young pupils in China. The same conclusion was drawn when AIDS and sex education in China were assessed in 2001[11]. To further emphasize that pupils in China are in need of more education, the study showed that the foreign medical students had received significantly more information before coming to university.

General Knowledge of HIV/AIDS

The knowledge regarding the distinction between HIV and AIDS was good. Three quarters of the students could correctly distinguish HIV from AIDS. This differs greatly from results in a similar study performed in Kazakhstan, where only 17.6% answered correctly [9].

There was a significant difference between all Chinese students and the foreign medical students in regard to what they believed to be the main route of transmission in China today. The majority of the Chinese students answered blood transfusion, while the foreign medical students almost exclusively answered sexual intercourse. These results can be explained by the fact that blood transfusions were once the main route of transmission in China, especially in Hubei province. In 1997, the Chinese government issued new laws and regulations, and therefore transmission *via* blood donations is now considered to be under control [12]. Sexual intercourse has just recently become the dominant mode of spread [2,5]. It can be assumed that the reason the foreign students answered sexual intercourse was that this is the main route of transmission throughout the world [13].

Most of the respondents knew the main routes of transmission. Compared to a study among Chinese college students in 2000, the awareness among the Wuhan students were better [14]. Still, about one third of the students had misconceptions concerning mosquitoes and kissing. The same misconceptions were found among students in Xinjiang [8]. Overall, the female students had better knowledge than the male students of the routes of transmission.

Almost eighty percent of the students answered that there are ways to protect oneself against sexually transmitted

HIV/AIDS. Most wrote in answer to the open corollary question: by condom use. This is in line with findings in a previous study at the same university where more than ninety percent knew that condom use prevents STDs [10].

An unexpected result was that the foreign medical students knew to a much larger extent than the Chinese medical students how to diagnose HIV. When asked to specify, most of the foreign students also gave more exact specifications, such as ELISA and WESTERN BLOT, while the Chinese students often wrote blood test or HIV test. This is surprising since the two groups study medicine at the same university.

More than eighty percent of the students knew that HIV/AIDS cannot be cured and that medication only slows down the progression. Looking at similar studies, the level of knowledge was higher among the Wuhan students [8,9]. In answer to the question of whether there were other treatments besides antiretroviral medicines, the Chinese medical students more often than the other student groups in the study answered that there were, referring to Chinese traditional medicines. There are not, to date, studies to provide sufficient information about the effects of Chinese traditional medicines on HIV [15].

Attitudes

Stigmatization of HIV/AIDS in China has been a major reason for the difficulties in reducing the rate of spread [1]. It has been shown that the degree of stigmatising beliefs is related to levels of income and education [16]. Among the Wuhan University students in the study, who are considered to be well educated, more than fifty percent answered either that they would or did not know if they would treat a person with HIV differently from others. This could be an indication that stigmatisation is still a substantial problem even among the well educated population. However, it must be taken into account that many students wrote that they would treat an infected person differently in a positive way.

Compared with students in Kazakhstan, the Wuhan students believed to a larger extent that there are special groups that are more likely to be infected with HIV than others. Prostitutes and drug addicts were pointed out in both studies [9]. This is in agreement with conclusions drawn when subpopulations of importance for HIV spread were investigated in 2005 [17].

Private Life

The vast majority of the students talk to their friends about general information regarding HIV/AIDS. It was much more common to talk to friends than parents, which is an indication that the subject is not openly discussed. However, more female than male students felt that they could talk to their parents. These findings are consistent with previous studies [8,14,18]. Chinese society is very traditional, and matters like HIV/AIDS are considered private. Fear and shame were considered reasons not to discuss subjects related to sexuality by many of the interviewed students.

It can be concluded that the Wuhan University students in the study did not have extensive sexual risk behavior.

Only 8.1% were sexually active, which is comparable with findings concerning other students in China [8,10,19]. The age of first sexual contact was also comparable [8,10]. Attitudes towards premarital sex have become more tolerant in the last few years in China [5,18]. In the interviews, students agreed that there has been a change of attitudes but that behavior has not yet changed.

A total of 67.5% of the students believed that both partners have equal responsibility for the use of condoms during sexual intercourse. This is twice as many as students in Xinjiang [8], probably due to cultural differences between the Chinese regions. Almost half of the students in the study wanted to take an HIV test. Many gave as a reason that they want to be healthy or just to be sure. This might be an indication of lack of knowledge with regard to the routes of transmission since it is not likely that they had been exposed to the virus. These results are also similar to the results concerning the Xinjiang students [8].

Future Epidemic in China

There is a great risk of a future epidemic in China according to the students. The main reasons for their opinion were the lack of knowledge in the general population and the change in sexual behavior among young people in China. Many believe that an epidemic can be prevented, but only if the government take more enforceable actions.

In conclusion, the Wuhan University students in the study had a positive and accepting attitude towards people living with HIV, and no extensive risk behavior. The level of knowledge varied between the three main groups and to some extent between genders. Overall, the level of knowledge was found to be moderate, and there is room for improvement. In China, there is a need for more sexual education both before and at university. It is especially important to increase the overall knowledge among the men and to give all increased awareness of the possibilities to prevent mother to child transmission of HIV and to reduce stigma by a deeper understanding on how HIV is not transmitted.

ACKNOWLEDGEMENTS

We would like to thank Professor Jing Sun for sharing her statistical knowledge and Professor Chunhong Wang for all her support.

To all the students at Professor Tan's laboratory and other friends, thank you for all the time and energy that you put into the study, and for making our stay in Wuhan unforgettable.

REFERENCES

- [1] Wu Z, Rou K, Cui H. The HIV/AIDS epidemic in China: history, current strategies and future challenges. *AIDS Educ Prevent* 2004;16(Suppl A): 7-17.
- [2] State Council AIDS Working Committee Office, UN Theme Group on AIDS in China: A Joint Assessment of HIV/AIDS Prevention, Treatment and Care in China. Available at: http://data.unaids.org/pub/Report/2008/china_2008_country_progress_report_en.pdf [Accessed April 7, 2008].
- [3] Zhang KL, Ma SJ. Epidemiology of HIV in China. Intravenous drug users, sexworkers, and large mobile populations are high risk groups. *BMJ* 2002; 324: 803-4.

- [4] Gorbacha PM, Ryan C, Saphonn V, Detels R. The impact of social, economic and political forces on emerging HIV epidemics. *AIDS* 2002; 16 (Suppl 4): S35-43.
- [5] Zhang K, Beck EJ. Changing sexual attitudes and behavior in China: Implications for the spread of HIV and other sexually transmitted diseases. *AIDS Care* 1999; 11(5): 581-9.
- [6] Médecins Sans Frontières. Available at: <http://www.msf.org.au/stories/twproject/2004/036twp.shtml> [Accessed April 8, 2008].
- [7] Wuhan University. Available at: <http://www.whu.edu.cn/en/about/index.html> [Accessed April 14, 2008].
- [8] Lönn E, Sahlholm K, Maimaiti R, Abdulkarim K, Andersson R. A traditional society in change encounters HIV/AIDS: knowledge, attitudes and risk behavior among students in Northwestern China. *AIDS Patient Care STDs* 2007; 21(1): 51-60.
- [9] Hansson L, Stockfelt L, Urazalin M, Andersson R, Ahlm C. HIV/AIDS awareness and risk behavior among students in Kazakhstan: a cross sectional survey. *BMC Int Health Hum Rights* 2008; 8: 14.
- [10] Tan X, Pan J, Zhou D, Wang C, Xie C. HIV/AIDS knowledge, attitudes and behaviors assessment of Chinese students: a questionnaire study. *Int J Environ Res Public Health* 2007; 4(3): 248-53.
- [11] Gao Y, Lu ZZ, Shi R, Sun XY, Cai Y. AIDS and sex education for young people in China. *Reprod Fertile Dev* 2001; 13: 729-37.
- [12] Wu Z, Rou K, Detels R. Prevalence of HIV infection among former commercial plasma donors in rural eastern China. *Health Policy Plan* 2001; 16(1): 41-6.
- [13] UNAIDS. 2006 report on the global AIDS epidemic: A UNAIDS 10th anniversary special edition. Available at: <http://www.unaids.org/en/KnowledgeCentre/HIVData/GlobalReport> [Accessed April 25, 2008].
- [14] Li X, Lin Ch, Gao Z, *et al.* HIV/AIDS knowledge and implications for health promotion programs among Chinese college students: geographic, gender and age differences. *Health Promot Int* 2004; 19 (3): 345-56.
- [15] Portsmouth S, Stebbing J, Keyi X, Jianping Z, Guohua P. HIV and AIDS in the Peoples Republic of China: a collaborative review. *Int J STD AIDS* 2003; (14): 757-61.
- [16] Liu H, Li X, Stanton B, *et al.* Relations of sexual risks and prevention practises with individuals' stigmatising beliefs towards HIV infected individuals: an exploratory study. *Sex Transm Infect* 2005; 6 (81): 511-6.
- [17] Qian ZH, Vermund SH, Wang N. Risk of HIV/AIDS in China: subpopulations of importance. *Sex Transm Infect* 2005; 81: 442-7.
- [18] Wang B, Li X, Stanton B, *et al.* Sexual attitudes, pattern of communication, and sexual behavior among unmarried out-of-school youth in China. *BMC Public Health* 2007; 7: 189.
- [19] Huang J, Bova C, Frennie KP, Rogers A, Williams AB. Knowledge, attitudes, behaviors, and perceptions of risk related to HIV/AIDS among Chinese university students in Hunan, China. *AIDS Patient Care STDS* 2005; 11(19): 769-77.

Received: April 24, 2009

Revised: July 21, 2009

Accepted: August 26, 2009

© Albrektsson *et al.*; Licensee *Bentham Open*.This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.